



**Roskilde
University**

A statistical mechanical approximation for the calculation of time auto-correlation functions

Dyre, Jeppe

Publication date:
1994

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Dyre, J. (1994). *A statistical mechanical approximation for the calculation of time auto-correlation functions*. Roskilde Universitet. Tekster fra IMFUFA No. 287 <http://milne.ruc.dk/lmfufaTekster/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 287

1994

**A Statistical Mechanical Approximation for the
Calculation of Time Auto-Correlation Functions**

By: Jeppe C. Dyre

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

ABSTRACT

This paper considers the problem of estimating the time auto-correlation function for a quantity that is defined in configuration space, given a knowledge of the mean-square displacement as function of time in this part of phase space. An approximate formula is derived which reduces the calculation of the time auto-correlation function to a "double canonical" average. In this approximation, the mean-square displacement itself may be evaluated from the "double partition function" in the case of Langevin dynamics. The scheme developed is illustrated by computer simulations of a simple one-dimensional systems, showing a good agreement between the exact time auto-correlation functions and those found by the approximation.

The calculation of a time auto-correlation function [1,2] is a straightforward matter in any computer simulation tracing the time evolution of a system [3-5]. However, computer simulations are not feasible today on time scales longer than microseconds. These time scales are relevant for viscous liquids approaching the glass transition. Therefore, one cannot by simulations calculate a number of experimentally accessible quantities in viscous liquids. Examples are the frequency-dependent viscosity [7], bulk modulus [8], dielectric constant [9], or specific heat [10,11], that all via the fluctuation-dissipation theorem [1,2,6] are given as Laplace transforms of a time auto-correlation function. In this situation one would like to have an approximate theory at hand. Focussing only on time auto-correlation functions of quantities $A(X)$ that are defined in configuration space, $X=(X_1, \dots, X_n)$, an approximation is proposed below, based on an ansatz for the joint probability of initial point X at $t=0$ and final point X' after time t , $P(X, X'; t)$. In terms of the joint probability the time auto-correlation function is given by

$$\langle A(0)A(t) \rangle = \int dX dX' A(X) A(X') P(X, X'; t) \quad (1)$$

If Z is the configurational partition function, given in terms of $\beta=1/(k_B T)$ and the potential energy $U(X)$ as

$$Z(\beta) = \int dX e^{-\beta U(X)} \quad (2)$$

and $G(X \rightarrow X'; t)$ is the Green's function, the exact expression for

the joint probability is

$$P(X, X'; t) = \frac{e^{-\beta U(X)}}{Z} G(X \rightarrow X'; t) \quad (3)$$

However, Eq. (3) is not very useful unless the Green's function is known. The principle of detailed balance implies that $P(X, X'; t) = P(X', X; t)$, a requirement any approximation should also satisfy to ensure time-reversal invariance.

The exact method for calculating $\langle A(0)A(t) \rangle$ is shown in Fig. 1 illustrating the path in configurational space. At a number of times t_1, \dots, t_n one computes the quantity $A(X(t_j))A(X(t_j+t))$, and $\langle A(0)A(t) \rangle$ is the average of this product as $n \rightarrow \infty$. Assuming here and henceforth that the X_i 's are simple rectangular coordinates and that $\langle A \rangle = 0$, one always finds $\langle A(0)A(t) \rangle \rightarrow 0$ as $t \rightarrow \infty$. This loss of correlation after long time comes about because the final point X' is far away from the initial point X . A measure of the distance travelled in time t is provided by the mean-square displacement, $\langle \Delta X^2(t) \rangle$: If $X \equiv X(t_j)$, $X' \equiv X(t_j+t)$, and $\langle \rangle$ denotes an average over j , the mean-square displacement is defined by

$$\langle \Delta X^2(t) \rangle = \langle (X - X')^2 \rangle \equiv \sum_{i=1}^n \langle (X_i - X'_i)^2 \rangle \quad (4)$$

Assuming that the mean-square displacement itself is a known function of time, the idea is now to estimate $\langle A(0)A(t) \rangle$ via

the "spatial" auto-correlation of A in configurational space evaluated at distances equal to $\sqrt{\langle \Delta X^2(t) \rangle}$. Before proceeding, we briefly discuss the physics of this way of thinking about the time auto-correlation function. A simple case is when the mean-square displacement is proportional to time (for $t \rightarrow \infty$ this is, of course, always the case). In this case, if the "spatial" correlation of A has a Gaussian distance decay, the time auto-correlation function is a simple exponential, corresponding to Debye relaxation. If, however, the spatial correlation of A has an exponential distance decay, the time auto-correlation function is a stretched exponential with exponent $1/2$. The latter case gives a reasonable fit to many experiments on viscous liquids [12]. The above picture of decomposing the time auto-correlation function into a) a "geometric" correlation and b) the distance travelled in a given time, is in harmony with another well-known property of viscous liquids. In these systems all linear relaxation functions have roughly the same average relaxation rate, a rate which slows down dramatically upon cooling. In the "geometric" picture, this is simply a consequence of the motion slowing down in configuration space, whereas the "spatial" correlation probably only change little upon cooling in a narrow range of temperatures. The mean-square displacement acts as a "molecular clock".

We now turn to the problem of estimating the joint probability $P(X, X'; t)$. In the thermodynamic limit $n \rightarrow \infty$ the relative fluctuations in the mean-square displacement go to zero,

and therefore the distance between $X=X(t_j)$ and $X'=X(t_j+t)$ is precisely $\sqrt{\Delta X^2(t)}$. Similarly, the relative fluctuations in potential energy go to zero, so the potential energy of both points X and X' is equal to $\langle U \rangle = -\frac{\partial \ln Z}{\partial \beta}$. The ansatz for

$P(X, X'; t)$ assumes equal probability for all pair of points with the correct distance and the correct potential energy. Thus,

$$P(X, X'; t) \propto \delta[(X-X')^2 - \langle \Delta X^2(t) \rangle] \delta[U(X) - \langle U \rangle] \delta[U(X') - \langle U \rangle] \quad (5)$$

In the thermodynamic limit there is "equipartition" between $U(X)$ and $U(X')$, and the last two delta functions may be replaced by a single delta function,

$$P(X, X'; t) \propto \delta[(X-X')^2 - \langle \Delta X^2(t) \rangle] \delta[U(X) + U(X') - 2\langle U \rangle] \quad (6)$$

The next step is to convert Eq. (6) to a "canonical" form, which is computationally more convenient than the "microcanonical" form. This is done by replacing the first delta function by $\exp[-a(X-X')^2]$ where a is a Lagrangian multiplier adjusted to give the correct mean-square displacement. Similarly, the second delta function is replaced by $\exp(-b[U(X) + U(X')])$ where b is adjusted to ensure that the average of $U(X) + U(X')$ is $2\langle U \rangle$. If the "double partition function"

$$D(a,b) = \int dx dx' e^{-a(X-X')^2 - b[U(X)+U(X')]} \quad (7)$$

is introduced, the final ansatz for the calculation of the time auto-correlation function is

$$\langle A(0)A(t) \rangle = \int \frac{dx dx'}{D(a,b)} A(X) A(X') e^{-a(X-X')^2 - b[U(X)+U(X')]} \quad (8)$$

In the thermodynamic limit Eq. (8) is equivalent to the average Eq. (1) over the distribution Eq. (5).

The two parameters a and b are determined in the following way. First, $b=b(a)$ is found from the condition that the average joint potential energy of initial and final point is $2\langle U \rangle$. Thus, $b(a)$ is determined from the condition that this average is independent of a :

$$\frac{d}{da} \frac{\partial \ln D}{\partial b} = 0 \quad (9)$$

Since $\frac{d}{da} = \partial_a + \frac{db}{da} \partial_b$ [with the standard abbreviated notation for partial derivatives], the expansion of Eq. (9) leads to the following first order differential equation for $b(a)$

$$\frac{db}{da} = \frac{\partial_a D \partial_b D - D \partial_{ab}^2 D}{D \partial_b^2 D - (\partial_b D)^2} \quad (10)$$

Once the function $b(a)$ has been determined, $a=a(t)$ is found from requiring the mean-square displacement calculated from $D(a,b(a))$ to be correct:

$$-\frac{\partial \ln D}{\partial a} = \langle \Delta X^2(t) \rangle \quad (11)$$

The short and long time limits are determined as follows. For $a(t)$ one clearly has

$$\begin{aligned} a(t=0) &= \infty \\ a(t=\infty) &= 0 \end{aligned} \quad (12)$$

In the limit of large times X and X' are far apart and $U(X)$ is uncorrelated with $U(X')$. In this limit $b=\beta$:

$$b(a=0) = \beta \quad (13)$$

In the short time limit the points X and X' are forced together. Thus, $P(X, X'; t) \propto \delta(X - X') \exp[-2bU(X)]$ for $t \rightarrow 0$ and Eq. (1) yields

$$\lim_{t \rightarrow 0} \langle A(0) A(t) \rangle = \frac{\int dX A^2(X) e^{-2bU(X)}}{\int dX e^{-2bU(X)}} \quad (14)$$

In order for this to give the correct canonical average one must have $b=\beta/2$, i. e.,

$$b(a=\infty) = \frac{\beta}{2} \quad (15)$$

The short time behavior of $a(t)$ may be derived directly from the equations of motion, as briefly sketched below. In the case of Newtonian dynamics, the Green's function at short times is easily found from the integrated equations of motion where the momentum is Gaussianly distributed (for simplicity it is assumed that all particles have the same mass m),

$$G(X \rightarrow X'; t) \propto \exp \left[-\frac{\beta}{2m} \sum_{i=1}^n \left(\frac{m}{t} (X'_i - X_i) + \frac{1}{2} \partial_i U t \right)^2 \right] . \quad (16)$$

To first order in t this yields

$$G(X \rightarrow X'; t) \propto \exp \left[-a(t) (X - X')^2 - \frac{\beta}{2} [U(X') - U(X)] \right] , \quad (17)$$

where

$$a(t) = \frac{\beta m}{2t^2} \quad (\text{Newtonian dynamics, } t \rightarrow 0) . \quad (18)$$

Note that via Eq. (3) this Green's functions confirms the form of the ansatz Eq. (8) for $t \rightarrow 0$, as well as the boundary condition Eq. (15). Next we consider the case of Langevin dynamics,

$$\dot{X}_i = -\mu \frac{\partial U}{\partial X_i} + \xi_i(t) , \quad (19)$$

with the standard Gaussian white noise term [14] $\langle \xi_i(t) \xi_j(t') \rangle = 2 \mu k_B T \delta_{i,j} \delta(t-t')$. From the equations of motion one finds that, because the integrated noise term is Gaussianly distributed,

$$G(X \rightarrow X'; t) \propto \exp \left[-\frac{\beta}{4\mu t} \sum_{i=1}^n (X'_i - X_i + \mu \partial_i U t)^2 \right] . \quad (20)$$

At short times this again leads to Eq. (17), where however now

$$a(t) = \frac{\beta}{4\mu t} \quad (\text{Langevin dynamics, } t \rightarrow 0) . \quad (21)$$

In the case of Langevin dynamics Eq. (8) may be applied to the calculation of the force-force time auto-correlation

function. This leads to an equation that in principle allows a calculation of $\langle \Delta X^2(t) \rangle$ directly from the double partition function. The mean-square displacement in time t is given by (sum over i)

$$\langle \Delta X^2(t) \rangle = \int_0^t dt' \int_0^t dt'' \langle \dot{X}_i(t') \dot{X}_i(t'') \rangle . \quad (22)$$

Since the noise terms are uncorrelated at different times, Eqs. (19) and (22) imply

$$\frac{d^2}{dt^2} \langle \Delta X^2(t) \rangle = 2 \langle \dot{X}_i(0) \dot{X}_i(t) \rangle = 2 \mu^2 \langle \partial_i U(0) \partial_i U(t) \rangle . \quad (23)$$

From Eq. (8) the force-force time auto-correlation function is rewritten as

$$\begin{aligned} \langle \partial_i U(0) \partial_i U(t) \rangle = \\ \frac{1}{b^2} \int \frac{dX dX'}{D(a, b)} [\partial_i e^{-bU(X)}] [\partial'_i e^{-bU(X')}] e^{-a(X-X')^2} \end{aligned} \quad (24)$$

By partial integrations one finds

$$\begin{aligned} \langle \partial_i U(0) \partial_i U(t) \rangle = \\ - 4 \frac{a^2}{b^2} \int \frac{dX dX'}{D(a, b)} (X-X')^2 e^{-a(X-X')^2 - b[U(X) + U(X')]} \\ = 4 \frac{a^2}{b^2} \frac{\partial \ln D}{\partial a} \end{aligned} \quad (25)$$

Thus, the equation for $a(t)$ is from Eqs. (11), (23) and (25)

$$\left(\frac{d^2}{dt^2} + 8 \mu^2 \frac{a^2}{b^2} \right) \frac{\partial \ln D}{\partial a} = 0 . \quad (26)$$

The expansion of Eq. (26) is straightforward, though tedious.

In order to check the validity of Eq. (8) a simple systems was studied numerically obeying Langevin dynamics. The system was chosen to be so simple that the integral in Eq. (8) may be evaluated "exactly", thus avoiding the noise of Monte Carlo simulations. No attempts were made to verify that Eq. (26) gives the correct $a(t)$. Instead the following procedure was followed. At a number of fixed a -values $b(a)$ was found from the requirement that the average joint potential energy is $2\langle U \rangle$. Then the mean-square displacement was evaluated for each a from Eq. (11) and also as function of time from the dynamical simulations, allowing an identification of the times corresponding to the fixed a -values. Finally, the time auto-correlation function was calculated from Eq. (8) at the fixed a -values. Figure 2 shows the results for $\langle X^3(0)X^3(t) \rangle$ for the Langevin motion of a particle in a double-well potential [14]. The full curve is the exact time auto-correlation function found by solving the Smoluchowski equation [13] and the dots give the prediction of Eq. (8). Results are shown for $\beta=2$ and for $\beta=8$ in dimensionless units.

In this paper a statistical mechanical approximation for the calculation of time auto-correlation functions was derived. The formalism assumes a knowledge of the mean-square displacement in configurational space as function of time; the mean-square displacement acts as the "molecular clock". The remaining "spatial" auto-correlation calculation is a "double canonical" average (Eq. (8)). Note that the corresponding double partition

function, $D(a,b)$, contains the ordinary configurational partition function as a special case, $Z^2(\beta) = D(0, \beta)$.

The approximation is only useful if $\langle \Delta X^2(t) \rangle$ is known. Experimentally, this quantity is accessible via the intermediate incoherent scattering function. In some cases a phenomenological estimate of the mean-square displacement may be given. Thus, for hopping in a rugged energy landscape where all minima are equal, the mean-square displacement is **universal** at low temperatures (except for trivial scalings), i.e., it is independent of the barrier height probability distribution. This has been shown recently [15] by effective medium calculations and by computer simulations of the frequency-dependent conductivity, which is simply related to the mean-square displacement [16]. Finally, there is the possibility that the mean-square displacement may be found approximately from Eq. (26) if Langevin dynamics is assumed, as is common, e. g., in polymer dynamics [17].

Equation (26) signals that Langevin dynamics plays a special role in the proposed scheme for calculation of time autocorrelation functions. A question of considerable interest is if and when Langevin dynamics can be expected to give the same time auto-correlation functions as Newtonian dynamics [18]. If the ansatz is correct, two different dynamics give the same time auto-correlation functions for any quantity defined in configuration space, if just the two dynamics give the same mean-square displacement. In this way the ansatz provides a mechanism for the consistency of any two types of dynamics.

ACKNOWLEDGEMENT

This work was supported by the Danish Natural Science Research Council.

REFERENCES

1. J. P. Boon and S. Yip, "Molecular Hydrodynamics" (McGraw Hill, New York, 1980).
2. J.-P. Hansen and I. R. Macdonald, "Theory of Simple Liquids", 2nd Ed. (Academic Press, New York, 1986).
3. M. P. Allen and D. J. Tildesley, "Computer Simulations of Liquids" (Clarendon Press, Oxford, 1987).
4. "Simulations of Liquids and Solids", Eds. G. Ciccotti, D. Frenkel, and I. R. Macdonald (North-Holland, Amsterdam, 1987).
5. "Computer Simulations in Chemical Physics", Eds. M. P. Allen and D. J. Tildesley (Kluwer Academic Publishers, Dordrecht, 1993).
6. R. Zwanzig, Ann. Rev. Phys. Chem. **16**, 67 (1965).
7. R. B. Bird, R. C. Armstrong, and O. Hassager, "Dynamics of Polymeric Liquids", 2nd Ed. (Wiley, New York, 1987), Vol. 1.
8. T. Christensen and N. B. Olsen, Phys. Rev. B **49**, 15396 (1994).

9. G. Williams, in: "Dielectric and Related Molecular Processes, Specialist Periodical Report, Vol. 2", Ed. M. Davies (Chem. Soc., London, 1975), p. 151.
10. T. Christensen, J. Phys. (Paris) Colloq. **46**, C8-635 (1985).
11. N. O. Birge and S. R. Nagel, Phys. Rev. Lett. **54**, 2674 (1985).
12. R. Böhmer, K. L. Ngai, C. A. Angell, and D. J. Plazek, J. Chem. Phys. **99**, 4201 (1993).
13. N. G. van Kampen, "Stochastic Processes in Physics and Chemistry" (Norht-Holland, Amsterdam, 1981).
14. M. Morillo and J. Gómez-Ordóñez, Phys. Rev. A **46**, 6738 (1992).
15. J. C. Dyre, Phys. Rev. B **49**, 11709 (1994); erratum Phys. Rev. B **50**, 9692 (1994).
16. H. Scher and M. Lax, Phys. Rev. B **7**, 4491 (1973).
17. M. Doi and S. F. Edwards, "The Theory of Polymer Dynamics" (Clarendon Press, Oxford, 1986).
18. H. Löwen, J.-P. Hansen, and J.-N. Roux, Phys. Rev. A **44**, 1169 (1991).

FIGURE CAPTIONS

Fig. 1:

Path in configuration space with coordinates $X=(X_1, \dots, X_n)$ illustrating the exact definition of the time auto-correlation function. At a number of times t_1, \dots, t_n one computes the quantity $A(X(t_j))A(X(t_j+t))$, and the time auto-correlation function is obtained as the average of this product as $n \rightarrow \infty$. In the thermodynamic limit the distance between the points $X(t_j)$ and $X(t_j+t)$ is the same for all j , because the relative distance fluctuations go to zero as the number of degrees of freedom go to infinity. This distance is the square root of the mean-square displacement in time t . In the approximation for evaluating the time auto-correlation function proposed here, equal probability is given to all pairs of initial ($t=0$) and final points after time t , that have the correct distance and where each point has the correct potential energy.

Fig. 2:

Log-log plot of $\langle X^3(0)X^3(t) \rangle$ as function of time for a Langevin particle in a double-well potential given in dimensionless units as $U(X) = (1/4)X^4 - (1/2)X^2$. The full curve is the exact time auto-correlation function evaluated by solving the Smoluchowski equation. The symbols give the predictions of Eq. (8); the system is so simple that no Monte Carlo simulation is necessary to evaluate Eq. (8). Results are shown for $\beta=2$ and for $\beta=8$.

Fig. 1





Fig. 2



Liste over tidligere udkomne tekster
tilsendes gerne. Henvendelse herom kan
ske til IMFUFA's sekretariat
tlf. 46 75 77 11 lokal 2263

217/92 "Two papers on APPLICATIONS AND MODELLING
IN THE MATHEMATICS CURRICULUM"

by: Mogens Niss

218/92 "A Three-Square Theorem"

by: Lars Kadison

219/92 "RUPNOK - stationær strømning i elastiske rør"

af: Anja Boisen, Karen Birkelund, Mette Olufsen

Vejleder: Jesper Larsen

220/92 "Automatisk diagnosticering i digitale kredsløb"

af: Bjørn Christensen, Ole Møller Nielsen

Vejleder: Stig Andur Pedersen

221/92 "A BUNDLE VALUED RADON TRANSFORM, WITH
APPLICATIONS TO INVARIANT WAVE EQUATIONS"

by: Thomas P. Branson, Gestur Ólafsson and
Henrik Schlichtkrull

222/92 On the Representations of some Infinite Dimensional
Groups and Algebras Related to Quantum Physics

by: Johnny T. Ottesen

223/92 THE FUNCTIONAL DETERMINANT

by: Thomas P. Branson

224/92 UNIVERSAL AC CONDUCTIVITY OF NON-METALLIC SOLIDS AT
LOW TEMPERATURES

by: Jeppe C. Dyre

225/92 "HATMODELLEN" Impedansspektroskopi i ultrarent
en-krystallinsk silicium

af: Anja Boisen, Anders Gorm Larsen, Jesper Varmer,
Johannes K. Nielsen, Kit R. Hansen, Peter Bøggild
og Thomas Hougaard

Vejleder: Petr Viscor

226/92 "METHODS AND MODELS FOR ESTIMATING THE GLOBAL
CIRCULATION OF SELECTED EMISSIONS FROM ENERGY
CONVERSION"

by: Bent Sørensen

227/92 "Computersimulering og fysik"

af: Per M.Hansen, Steffen Holm,
Peter Maibom, Mads K. Dall Petersen,
Pernille Postgaard, Thomas B.Schrøder,
Ivar P. Zeck

Vejleder: Peder Voetmann Christiansen

228/92 "Teknologi og historie"

Fire artikler af:

Mogens Niss, Jens Høyrup, Ib Thiersen,
Hans Hedal

229/92 "Masser af information uden betydning"

En diskussion af informationsteorien
i Tor Nørretranders' "Mærk Verden" og
en skitse til et alternativ baseret
på andenordens kybernetik og semiotik.

af: Søren Brier

230/92 "Vinklens tredeling - et klassisk
problem"

et matematisk projekt af

Karen Birkelund, Bjørn Christensen

Vejleder: Johnny Ottesen

231A/92 "Elektrondiffusion i silicium - en
matematisk model"

af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen

Vejledere: Johnny Ottesen, H.B.Hansen

231B/92 "Elektrondiffusion i silicium - en
matematisk model" Kildetekster

af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen

Vejledere: Johnny Ottesen, H.B.Hansen

232/92 "Undersøgelse om den simultane opdagelse
af energiens bevarelse og isærdeles om
de af Mayer, Colding, Joule og Helmholtz
udførte arbejder"

af: L.Arleth, G.I.Dybkjær, M.T.Østergård

Vejleder: Dorthe Posselt

233/92 "The effect of age-dependent host
mortality on the dynamics of an endemic
disease and
Instability in an SIR-model with age-
dependent susceptibility

by: Viggo Andreassen

234/92 "THE FUNCTIONAL DETERMINANT OF A FOUR-DIMENSIONAL
BOUNDARY VALUE PROBLEM"

by: Thomas P. Branson and Peter B. Gilkey

235/92 OVERFLADESTRUKTUR OG POREUDVIKLING AF KOKS

- Modul 3 fysik projekt -

af: Thomas Jessen

- 236a/93 INTRODUKTION TIL KVANTE
HALL EFFEKTEN
af: Anja Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 236b/93 STRØMSSAMMENBRUD AF KVANTE
HALL EFFEKTEN
af: Anja Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 237/93 The Wedderburn principal theorem and
Shukla cohomology
af: Lars Kadison
- 238/93 SEMIOTIK OG SYSTEMEGENSKABER (2)
Vektorbånd og tensorer
af: Peder Voetmann Christiansen
- 239/93 Valgsystemer - Modelbygning og analyse
Matematik 2. modul
af: Charlotte Gjerrild, Jane Hansen,
Maria Hermannsson, Allan Jørgensen,
Ragna Clauson-Kaas, Poul Lützen
Vejleder: Mogens Niss
- 240/93 Patologiske eksempler.
Om sære matematiske fisks betydning for
den matematiske udvikling
af: Claus Dræby, Jørn Skov Hansen, Runa
Ulsøe Johansen, Peter Meibom, Johannes
Kristoffer Nielsen
Vejleder: Mogens Niss
- 241/93 FOTOVOLTAISK STATUSNOTAT 1
af: Bent Sørensen
- 242/93 Brovedligeholdelse - bevar mig vel
Analyse af Vejdirektoratets model for
optimering af broreparationer
af: Linda Kyndlev, Kare Fundal, Kamma
Tulinius, Ivar Zeck
Vejleder: Jesper Larsen
- 243/93 TANKEEKSPERIMENTER I FYSIKKEN
Et 1.modul fysikprojekt
af: Karen Birkelund, Stine Sofia Korremann
Vejleder: Dorthe Posselt
- 244/93 RADONTRANSFORMATIONEN og dens anvendelse
i CT-scanning
Projektrapport
af: Trine Andreasen, Tine Guldager Christiansen,
Nina Skov Hansen og Christine Iversen
Vejledere: Gestur Olafsson og Jesper Larsen
- 245a+b
/93 Time-Of-Flight målinger på krystallinske
halvledere
Specialerapport
af: Linda Szkotak Jensen og Lise Odgaard Gade
Vejledere: Petr Viscor og Niels Boye Olsen
- 246/93 HVERDAGSVIDEN OG MATEMATIK
- LÆREPROCESSER I SKOLEN
af: Lena Lindenskov, Statens Humanistiske
Forskningsråd, RUC, IMFUFA
- 247/93 UNIVERSAL LOW TEMPERATURE AC CON-
DUCTIVITY OF MACROSCOPICALLY
DISORDERED NON-METALS
by: Jeppe C. Dyre
- 248/93 DIRAC OPERATORS AND MANIFOLDS WITH
BOUNDARY
by: B. Booss-Bavnbek, K.P.Wojciechowski
- 249/93 Perspectives on Teichmüller and the
Jahresbericht Addendum to Schappacher,
Scholz, et al.
by: B. Booss-Bavnbek
With comments by W.Abikoff, L.Ahlfors,
J.Cerf, P.J.Davis, W.Fuchs, F.P.Gardiner,
J.Jost, J.-P.Kahane, R.Lohan, L.Lorch,
J.Radkau and T.Söderqvist
- 250/93 EULER OG BOLZANO - MATEMATISK ANALYSE SET I ET
VIDENSKABSTEORETISK PERSPEKTIV
Projektrapport af: Anja Juul, Lone Michelsen,
Tomas Højgård Jensen
Vejleder: Stig Andur Pedersen
- 251/93 Genotypic Proportions in Hybrid Zones
by: Freddy Bugge Christiansen, Viggo Andreasen
and Ebbe Thue Poulsen
- 252/93 MODELLERING AF TILFÆLDIGE FÆNOMENER
Projektrapport af: Birthe Friis, Lisbeth Helmgård,
Kristina Charlotte Jakobsen, Marina Mosbæk
Johannessen, Lotte Ludvigsen, Mette Haas Nielsen
- 253/93 Kuglepakning
Teori og model
af: Lise Arleth, Kåre Fundal, Nils Kruse
Vejleder: Mogens Niss
- 254/93 Regressionsanalyse
Materiale til et statistikkursus
af: Jørgen Larsen
- 255/93 TID & BETINGET UAFHÆNGIGHED
af: Peter Harremoës
- 256/93 Determination of the Frequency Dependent
Bulk Modulus of Liquids Using a Piezo-
electric Spherical Shell (Preprint)
by: T. Christensen and N.B.Olsen
- 257/93 Modellering af dispersion i piezoelektriske
keramikker
af: Pernille Postgaard, Jannik Rasmussen,
Christina Specht, Mikko Østergård
Vejleder: Tage Christensen
- 258/93 Supplerende kursusedmateriale til
"Lineære strukturer fra algebra og analyse"
af: Mogens Brun Heefelt
- 259/93 STUDIES OF AC HOPPING CONDUCTION AT LOW
TEMPERATURES
by: Jeppe C. Dyre
- 260/93 PARTITIONED MANIFOLDS AND INVARIANTS IN
DIMENSIONS 2, 3, AND 4
by: B. Booss-Bavnbek, K.P.Wojciechowski

- 261/93 OPGAVESAMLING
Bredde-kursus i Fysik
Eksamensopgaver fra 1976-93
- 262/93 Separability and the Jones Polynomial
by: Lars Kadison
- 263/93 Supplerende kursusmateriale til "Lineære strukturer fra algebra og analyse" II
af: Mogens Brun, Heefelt
- 264/93 FOTOVOLTAISK STATUSNOTAT 2
af: Bent Sørensen
-
- 265/94 SPHERICAL FUNCTIONS ON ORDERED SYMMETRIC SPACES
To Sigurdur Helgason on his sixtyfifth birthday
by: Jacques Faraut, Joachim Hilgert and Gestur Olafsson
- 266/94 Kommensurabilitets-oscillationer i laterale supergitre
Fysikspeciale af: Anja Boisen, Peter Bøggild, Karen Birkelund
Vejledere: Rafael Taboryski, Poul Erik Lindelof, Peder Voetmann Christiansen
- 267/94 Kom til kort med matematik på Eksperimentarium - Et forslag til en opstilling
af: Charlotte Gjerrild, Jane Hansen
Vejleder: Bernhelm Booss-Bavnbek
- 268/94 Life is like a sewer ...
Et projekt om modellering af aorta via en model for strømning i kloakrør
af: Anders Marcussen, Anne C. Nilsson, Lone Michelsen, Per M. Hansen
Vejleder: Jesper Larsen
- 269/94 Dimensionsanalyse en introduktion metaprojekt, fysik
af: Tine Guldager Christiansen, Ken Andersen, Nikolaj Hermann, Jannik Rasmussen
Vejleder: Jens Højgaard Jensen
- 270/94 THE IMAGE OF THE ENVELOPING ALGEBRA AND IRREDUCIBILITY OF INDUCED REPRESENTATIONS OF EXPONENTIAL LIE GROUPS
by: Jacob Jacobsen
- 271/94 Matematikken i Fysikken.
Opdaget eller opfundet
NAT-BAS-projekt
vejleder: Jens Højgaard Jensen
- 272/94 Tradition og fornyelse
Det praktiske elevarbejde i gymnasiets fysikundervisning, 1907-1988
af: Kristian Hoppe og Jeppe Guldager
Vejledning: Karin Beyer og Nils Hybel
- 273/94 Model for kort- og mellemdistanceløb
Verifikation af model
af: Lise Fabricius Christensen, Helle Pilemann, Bettina Sørensen
Vejleder: Mette Olufsen
- 274/94 MODEL 10 - en matematisk model af intravenøse anæstetikas farmakokinetik
3. modul matematik, forår 1994
af: Trine Andreasen, Bjørn Christensen, Christine Green, Anja Skjoldborg Hansen, Lisbeth Helmggaard
Vejledere: Viggo Andreasen & Jesper Larsen
- 275/94 Perspectives on Teichmüller and the Jahresbericht 2nd Edition
by: Bernhelm Booss-Bavnbek
- 276/94 Dispersionsmodellering
Projektrapport 1. modul
af: Gitte Andersen, Rehannah Borup, Lisbeth Friis, Per Gregersen, Kristina Vejro
Vejleder: Bernhelm Booss-Bavnbek
- 277/94 PROJEKTARBEJDSPEDAGOGIK - Om tre tolkninger af problemorienteret projektarbejde
af: Claus Flensted Behrens, Frederik Voetmann Christiansen, Jørn Skov Hansen, Thomas Thingstrup
Vejleder: Jens Højgaard Jensen
- 278/94 The Models Underlying the Anaesthesia Simulator Sophus
by: Mette Olufsen(Math-Tech), Finn Nielsen (RISØ National Laboratory), Per Føge Jensen (Herlev University Hospital), Stig Andur Pedersen (Roskilde University)
- 279/94 Description of a method of measuring the shear modulus of supercooled liquids and a comparison of their thermal and mechanical response functions.
af: Tage Christensen
- 280/94 A Course in Projective Geometry
by Lars Kadison and Matthias T. Kromann
- 281/94 Modellering af Det Kardiovaskulære System med Neural Puls kontrol
Projektrapport udarbejdet af:
Stefan Frello, Runa Ulsøe Johansen, Michael Poul Curt Hansen, Klaus Dahl Jensen
Vejleder: Viggo Andreasen
- 282/94 Parallell algoritmer
af: Erwin Dan Nielsen, Jan Danielsen, Niels Bo Johansen

- 283/94 Grænser for tilfældighed
(en kaotisk talgenerator)
af: Erwin Dan Nielsen og Niels Bo Johansen
- 284/94 Det er ikke til at se det, hvis man ikke
lige ve' det!
Gymnasiematematikens begrundelsesproblem
En specialerapport af Peter Hauge Jensen
og Linda Kyndlev
Vejleder: Mogens Niss
- 285/94 Slow coevolution of a viral pathogen and
its diploid host
by: Viggo Andreassen and
Freddy B. Christiansen
- 286/94 The energy master equation: A low-temperature
approximation to Bässler's random walk model
by: Jeppe C. Dyre